Energy Multiple Choice Homework

PSI Physics

Name

Multiple Choice Questions



- 1. A block of mass m is pulled over a distance d by an applied force F which is directed in parallel to the displacement. How much work is done on the block by the force F?
 - A. mFd
 - B. zero
 - C. Fd
 - D.
 - E. -Fd



2. A block of mass m is moved over a distance d. An applied force F is directed perpendicularly to the block's displacement. How much work is done on the block by the force F?

A. mFd

- B. zero
- C. Fd
- D.
- E.-Fd



- 3. A block of mass m is moved over a distance d. An applied force F is opposite to the block's displacement. How much work is done on the block by the force F?
 - A. mFd
 - B. zero
 - C. Fd
 - D.
 - E.-Fd



- 4. A truck driver is trying to push a loaded truck with an applied force. Unfortunately, his attempt was unsuccessful the truck stays stationary no matter how hard the driver pushes. How much work is done by the driver?
 - A. Fd B. -Fd C. D. E. Zero



5. A spacecraft moves around Earth in a circular orbit with a constant radius. How much work is done by the gravitational force on the spacecraft during one revolution?

A. $F_G d$ B. - $F_G d$ C. mgh D. $\frac{1}{2} mv^2$ E. zero



6. A construction worker holds a heavy tool box. How much work is done by the worker?

A. F_Gd

- B. F_Gd
- C. mgh
- D. ½ mv²
- E. zero
- 7. A container with a mass of 5 kg is lifted to a height of 8 m. How much work is done by the gravitational force?

A. 400 J B. -400 J C. zero D. 50 J E. -50J

8. A container with a mass of 5 kg is lifted to a height of 8 m. How much work is done by the external force?

A. 400 J B. -400 J C. zero D. 50 J E. -50J

- 9. A container with a mass of 5 kg is lifted to a height of 8 m and then returned back to the ground level. How much work is done by the gravitational force?
 - A. 400 J B. -400 J C. zero D. 50 J E. -50J
- 10. An object is thrown straight up. Which of the following is true about the sign of work done by the gravitational force while the object moves up and then down?
 - A. Work is positive on the way up, work is positive on the way down
 - B. Work is negative on the way up, work is negative on the way down
 - C. Work is negative on the way up, work is positive on the way down
 - D. Work is positive on the way up, work is negative on the way down
 - E. Work is zero the way up, work is zero on the way down



11. The force as a function of displacement of a moving object is presented by the graph. How much work is done when the object moves from 0 m to 8 m?

A. 40 J B. 20 J	C. 0 J	D. 10 J	E. 5 J
-----------------	--------	---------	--------



12. The force as a function of displacement of a moving object is presented by the graph. How much work is done when the object moves from 0 m to 5 m?





13. The force as a function of displacement of a moving object is presented by the graph. How much work is done when the object moves from 5 m to 8 m?



14. The force as a function of displacement of a moving object is presented by the graph. How much work is done when the object moves from 0 m to 8 m?

A. 30 J B. 15 J C. 18 J D. 9 J E. 20 J



- 15. An applied force F accelerates an object from rest to a velocity v. How much work is done by the applied force F?
 - A. ½ mv²
 - B. mgh
 - C. $\frac{1}{2}$ kx²
 - D. mFd
 - E. Zero
- 16. What happens to the kinetic energy of a moving object if the net work done is positive?
 - A. The kinetic energy increases
 - B. The kinetic energy decreases
 - C. The kinetic energy remains the same
 - D. The kinetic energy is zero
 - E. The kinetic energy becomes negative



17. A block of mass m = 50 kg moves on a rough horizontal surface with a coefficient of kinetic friction μ = 0.5. The traveled distance is 20 m. How much work is done by the friction force?

A. 1000 J B. 2000 J C. 3000 J D. 4000 J E. -5000 J

- 18. What happens to the gravitational potential energy of a moving object if the work done by the gravitational force is negative?
 - A. The potential energy increases
 - B. The potential energy decreases
 - C. The potential energy remains the same
 - D. The potential energy is zero
 - E. The potential energy becomes negative



- 19. An object I with a mass of 4 kg is lifted vertically 3 m from the ground level; another object II with a mass of 2 kg is lifted 6 m up. Which of the following statements is true?
 - I. Object I has greater potential energy since it is heavier
 - II. Object II has greater potential energy since it is lifted to a higher position
 - III. Two objects have the same potential energy

	A. I	B. II	C. III	D. I and II	E. II and III
--	------	-------	--------	-------------	---------------



20. A 4 kg block is attached to a vertical spring with a spring constant 800 N/m. The spring stretches 5 cm down. How much elastic potential energy is stored in the system?

A. 1.0 J B. 0.5 J C. 1.5 J D. 2.0 J E. 2.5 J



21. A heavy block is suspended from a vertical spring. The elastic potential energy is stored in the spring is 2 J. What is the spring constant if the elongation of the spring is 10 cm?

A. 400 N/m B. 300 N/m C. 200 N/m D. 100 N/m E. 50 N/m



22. A heavy block is suspended from a vertical spring. The elastic potential energy is stored in the spring is 0.8 J. What is the elongation of the spring if the spring constant is 100 N/m?

A, Z CIII D, 4 CIII C, 0 CIII D, 11 CIII L, 13 CIII	A. 2 cm	B. 4 cm	C. 8 cm	D. 11 cm	E. 13 cm
---	---------	---------	---------	----------	----------



23. The elastic force as a function of displacement presented by the graph. How much elastic potential energy is stored in the spring when it is stretched by 10 cm?



24. A bullet penetrates a wooden block and loses its velocity by a half. What is the ratio between the initial kinetic energy of the bullet and kinetic energy when the bullet leaves the block?

$$A. = B. = C. = D. = E. =$$

25. A truck drives slams on the brakes of a moving truck with a constant velocity v, as a result of his action the truck stops after traveling a distance d. If the driver had been traveling with twice the velocity, what would be the stopping distance compared to the distance in the first trial?

A. Two times greater

- B. Four times greater
- C. The same
- D. Half as much
- E. One-quarter as much
- 26. What happens to the total energy of a moving object if all the applied forces are conserved?
 - A. It increases
 - B. It decreases
 - C. It remains constant
 - D. The velocity is required to answer this question
 - E. The altitude is required to answer this question
- 27. A machine does 2500 J of work in 1 min. What is the power developed by the machine?

A. 21 W B. 42 W C. 150 W D. 2500 W E. 150000 W

- 28. A car travels with a constant speed of 15 m/s. The car's engine produces a 4000 N pushing force in order to keep the speed constant. How much power is developed by the engine?
 - A. 60 W B. 600 W C. 6000 W D. 60000 W E. 600000W



29. A block with a mass of m crosses a rough horizontal surface at a constant speed of v. The coefficient of kinetic friction between the block and the surface is μ . How much power must be produced in order to overcome the friction force?

A. mg B. µmg C. zero D. µg E. µmgv

30. A motorbike engine can develop a power of 90000 W in order to keep a constant velocity of 30 m/s. What is the pushing force?

A. 3000 N B. 30000 N C. 300000 N D. 300 N E. 30 N

Answers

- 1. C 2. B 3. E 4. E 5. E 6. E 7. B 8. A 9. C 10. C 11. A 12. B 13. D 14. E 15. A 16. A 17. E 18. A 19. C 20. A 21. A 22. E 23. C 24. D 25. B 26. C 27. B 28. D
- 29. E
- 30. A